

PATIENT NAME: [patient name]

EXAM DATE: [exam date] **AGE:** [age] **SEX:** [sex] **I.D.#:** [patient I.D.]

CT URINARY TRACT WO	Kidney stones, flank pain
FORMS	Consent Form Pregnancy Status Form (female patients)
ORAL CONTRAST	NONE 1 GLASS OF WATER ON TABLE, BEFORE SCANNING
IV CONTRAST	NONE UNLESS PHYS ORDERS WITH CONTRAST
POSITIONING	SUPINE; FEET FIRST; ARMS UP & OUT, CENTER AT XYPHOID INSPIRATION BREATH-HOLD
SCOUT	S60/I 450, 120 KVP @ 10 MA, AP & LAT
TECHNIQUE – 16 & 64 SLICE	HELICAL; 2.5 MM, STD W400/L40 KVP 120 @ AUTO/SMART MA-40% ASIR SFOV - LG BODY/DFOV-36 CM (FIT TO ANATOMY) SCAN FROM DIAPHRAGM DOWN THROUGH BLADDER
TECHNIQUE - 128 SLICE	
RECONS	16 SLICE-1.25 MM/INT 0.625 MM, STD AT W400/L40 64 SLICE-0.625 MM, STD AT W400/L40 (DMPR) 2.5 MM, LUNG AT W1400/L-600
REFORMATIONS	CORONALS AVG 2.5/2.5, W400/L40
PACS	SCOUT 2.5 MM -UT ABD COR & ABD SAG
CHARGE	URINARY TRACT W/O
REMARKS	USE LOW DOSE PROTOCOL (MA 125), IF PT HAS HAD PREVIOUS SCAN WITHIN ONE MONTH

CT PREPS	All CT contrast patients should be NPO for 4 hours before their exam. All patients should drink plenty of water before their exam
Routine Abdomen and Pelvis	For: Diverticulitis, pain, pancreatitis, appendicitis, elevated labs or cancer (not for adrenal, liver, GIST, stromal, or pancreas cancer). Outpatients: Patients are to receive one 50 ml bottle of Omni 140, consent form, and prep sheet. Patient to mix Omni 140 50 cc into 32 oz water.

